

Accessibility

Patient information leaflets are available on our website:
www.oxfordhealth.nhs.uk/leaflets

Get in touch

Address Oxford Health NHS Foundation Trust
Trust Headquarters
Littlemore Mental Health Centre
Sandford Road, Oxford OX4 4XN

Phone 01865 901 000
Email enquiries@oxfordhealth.nhs.uk
Website www.oxfordhealth.nhs.uk

Feedback

Our Patient Advice and Liaison Service (PALS) provides advice and support to patients, families, and carers, helping to resolve any problems, concerns, or complaints you may have.

Phone 0800 328 7971
Email PALS@oxfordhealth.nhs.uk

Become a member of our foundation trust: www.ohftnhs.uk/membership

Support Oxford Health Charity, making a difference to patients, their families and staff who care for them: www.oxfordhealth.charity



Bladder and Bowel Service

Men's guide to the pelvic floor muscles

Contents

Introduction.....	3
What do they do?.....	3
Factors leading to incontinence or weakness.....	4
How to exercise your pelvic floor muscles.....	5
Exercise programme.....	6
Personalised programme/advice from your physiotherapist.....	8
Further useful tips and advice	10
Bladder emptying.....	12
Your appointment:.....	14
Accessibility	16
Get in touch.....	16
Feedback	16

Your appointment:

Date

Time






Please contact the number below if you are unable to come to this appointment, giving as much notice as possible.

The information contained in this leaflet is intended for educational use only and not for the diagnosis or treatment of a specific condition, which should only be undertaken by a qualified healthcare professional.

Introduction

Many men suffer from weakness of their pelvic floor muscles.



Common symptoms include:

-  leaking urine with activity, for example during running,
-  jumping, coughing and sneezing
-  a sudden and urgent need to pass urine
-  leakage of stool from the back passage
-  erectile dysfunction (the pelvic floor helps you to maintain erections).


This information is a guide to be used alongside the advice and guidance given by your Physiotherapist.

What do they do?

The pelvic floor muscles are a supportive sling of muscles, stretching from the tailbone at the back to the pubic bone at the front. They are responsible for:

-  supporting the pelvic organs - the bladder and bowel, especially when standing; they also help protect these organs from external damage
-  a role in erectile function during intercourse.

Pelvic Floor Muscles



Your pelvic floor muscles contract when you cough, sneeze, or laugh, preventing leakage of urine.

They are kept under slight tension so that when you pass urine or open your bowels the muscles should relax and then tighten afterwards. This prevents leakage and controls the passing of urine, bowel, and gas motions.

Factors leading to incontinence or weakness

- i** prostate surgery
- i** long term cough
- i** constipation/straining to empty your bowels
- i** being overweight

Holding too long can result in the bladder not working correctly in the long term as the bladder muscle can become overstretched.

If problems continue you may need to seek further medical advice.

Bladder emptying

It is important to empty your bladder completely each time you go to the toilet. Take your time. If you think your bladder may not be completely empty, try rocking your pelvis forwards and backwards; it may help to empty it a little more.

Frequency of bladder emptying

The bladder normally empties between five and eight times per day. This usually means emptying your bladder every two to five hours. Leaving too little or too long a time between emptying may upset this normal function.

Less than two hours or more than five hours may lead to problems. You may need to train your bladder to hold more by trying delaying techniques when you feel the urge to go, such as:

- i** keep calm and tighten your pelvic floor muscles
- i** sit on a hard surface
- i** curl your toes hard
- i** distract your mind by counting backwards, singing a song, etc.

- i** pelvic trauma or some surgery
- i** repeated heavy lifting.

Pelvic floor muscles can be strengthened just like any other weak or damaged muscles. They need to be regularly exercised to work well.

How to exercise your pelvic floor muscles

- ✓** Sit, stand or lie comfortably with your knees slightly apart.
- ✓** Imagine you are trying to stop yourself from passing wind at the same time as if you are trying to stop passing urine; slowly squeeze and lift the muscles. You may feel the base of your penis move slightly up towards your abdomen.
- ✓** A feeling of gentle tightening in your lower abdomen is normal. Try to avoid pulling in your stomach, squeezing your legs together, tightening your buttocks or holding your breath. This will help to ensure that only your pelvic floor muscles are working.
- ✓** NUTS to GUTS is easy to remember.

Exercise programme

First you need to find your starting point.

To do this, tighten your pelvic floor muscles and hold the contraction for as many seconds as you can, up to ten seconds.

How long can you hold the contraction?

..... seconds.

Release the contraction and rest for the same time. Repeat the tightening, hold, release, as many times as you can, up to a maximum of ten.

How many times can you repeat the contraction?

..... times.

For example, hold two seconds, rest two seconds, repeat four times.

These exercises will build the endurance of the muscles (the muscles can work harder for longer). As you improve, you will notice that you can hold your contraction for longer and do more repetitions, compared to your starting point.

When laughing, coughing, and sneezing, your muscles need to be able to react quickly. It is also important to practise quick contractions. To do this, tighten the muscles quickly and strongly and then relax.

4. Exercises to avoid if you think your pelvic floor is weak

- ❗ double leg lifts
- ❗ straight leg sit-ups
- ❗ repetitive heavy lifting
- ❗ high impact exercise such as jumping and running.

These activities put increased stress and pressure on the pelvic floor muscles.

5. Movement changes

Try to tighten your pelvic floor muscles before coughing, sneezing, and laughing. This will help to prevent leakage.

Further useful tips and advice

1. Avoid being overweight

Maintaining the correct weight can make a big difference to your symptoms, by reducing the strain on the muscles that help to control bladder and bowel movements.

2. Avoid constipation

Straining to open your bowels stretches the pelvic floor muscles, making them weaker. It is important to drink plenty of fluids and you may need to adjust your diet. This is particularly important if you have just had prostate surgery.

3. Drink enough fluids

Reduce your caffeine intake, as these drinks may irritate your bladder. Try to drink more water and aim to drink between 3 - 4 pints (1.5 - 2 litres) of fluid a day.

Do not restrict your fluid intake. This can make your symptoms worse as your urine will be more concentrated. Regular drinking at the same times each day is best.

How many quick contractions can you do?

..... times.

Aim for ten repetitions.

Aim to increase the number of repetitions of slow and quick exercises over the next six months so that you can do up to ten at a time. You will need to practise three times per day.

Personalised programme/advice from your physiotherapist

1. Timed contractions

Pull up and hold as tightly as you can for seconds,
rest for seconds,
repeat times
do this times a day.

2. Fast contractions

Repeat times and do times a day.

The most difficult part of the programme is remembering to do the exercises. The tips below may help to trigger your memory:

- ✓ Put coloured sticky notes or dots in places that will catch your eye (on the fridge, the kettle or your watch).
- ✓ Set an alarm on your watch/phone.
- ✓ Exercise after every time you empty your bladder.
- ✓ Exercise when waiting for the kettle to boil or when washing your hands.

Personalised illustration box (for use by your therapist)